arXiv:math/0504476v1 [math.CA] 22 Apr 2005

Some remarks on a paper by L. Carlitz

Diego Dominici [*]
Department of Mathematics
State University of New York at New Paltz
75 S. Manheim Blvd. Suite 9
New Paltz, NY 12561-2443
USA
Phone: (845) 257-2607
Fax: (845) 257-3571

Abstract

We study a family of orthogonal polynomials which generalizes a sequence of polynomials considered by L. Carlitz. We show that they are a special case of the Sheffer polynomials and point out some interesting connections with certain Sobolev orthogonal polynomials.

Keywords: Sheffer polynomials, Favard theorem, Sobolev orthogonal polynomials, generating functions, hypergeometric polynomials, special functions.

MSC-class: 33A65 (Primary) 11B83, 46E39 (Secondary)

1 Introduction

In [32] R. Kelisky defined a set of rational integers T_n by means of the exponential generating function

$$\exp\left[\arctan\left(z\right)\right] = \sum_{n=0}^{\infty} T_n \frac{z^n}{n!}.$$

---

[*] e-mail: dominicd@newpaltz.edu

L. Carlitz extended Kelisky's idea by defining the polynomials $T_n(x)$, with

$$\exp\left[x \arctan\left(z\right)\right] = \sum_{n=0}^{\infty} T_n(x) \frac{z^n}{n!}. \tag{1}$$

In his article [15], he showed that $T_n(x)$ satisfies the recurrence

$$T_{n+1} = xT_n - n(n-1)T_{n-1}, \quad n \geq 0, \tag{2}$$

where $T_{-1}(x) = 0$ and $T_0(x) = 1$. He also proved [16] several arithmetic properties of $T_n(x)$.

In [31] S. Kaijser considered the recurrence (2) and concluded that:

1. The function $T_n(x)$ is a monic polynomial of degree n.
2. The set $\left\{\frac{1}{n!}T_n(x) \mid n \geq 0\right\}$ is an orthonormal basis of the Hilbert space $H^2(S, P)$, where $S = \{z \in \mathbb{C} \mid \operatorname{Im}(z) \in [-1, 1]\}$ and P is the Poisson measure for 0.
3. The norm of the polynomial $\frac{1}{n!}T_n(x)$ is $\sqrt{2}$ if $n \geq 1$ and 1 if $n = 0$.
4. The exponential generating function of $T_n(x)$ is $\exp\left[x \arctan\left(z\right)\right]$.

The same results and some extensions were presented by T. Araaya in [6]. He also showed that

$$\frac{1}{n!}T_n(x) = \frac{x}{n} P_{n-1}^{(1)}\left(\frac{x}{2}; \frac{\pi}{2}\right), \quad n \geq 1 \tag{3}$$

where $P_n^{(\lambda)}\left(x; \phi\right)$ is the Meixner-Pollaczek polynomial [17].

In this paper we extend (1) by consider the following problem.

Problem 1 *For which functions $f(z)$, will the polynomials $\Psi_n(x)$, generated by*

$$\exp\left[xf\left(z\right)\right] = \sum_{n=0}^{\infty} \Psi_n(x) \frac{z^n}{n!}, \tag{4}$$

form an orthogonal set?

In Section 2 we show that the, somewhat surprising, answer is that the function $f(z)$ must be of the $\arctan(\cdot)$ type.

Problem 1 is a particular case of the Sheffer problem [57] of characterizing the orthogonal polynomials $S_n(x)$ generated by

$$F(z)\exp\left[xf\left(z\right)\right]=\sum_{n=0}^{\infty}\Psi_n(x)\frac{z^n}{n!}$$

where $F(0)=1$ and $f(0)=0$. Although the Sheffer polynomials have been studied extensively [3], [10], [19], [20], [26], [55] [56], [58], the limiting case (4) with $F(z)=1$ does not appear to have been considered before.

It is known [2], [17], that the Sheffer polynomials reduce to the Hermite, Laguerre, Charlier, Meixner or Meixner-Pollaczek polynomials, depending on the choice of $F(z)$ and $f\left(z\right)$. Motivated by (3), we wondered if some of the polynomials generated by (4) will form a new class. In Section 3 we show that they will be limiting cases of the Laguerre, Meixner-Pollaczek or Meixner polynomials.

An interesting property of these polynomials is that they do not form an orthogonal set with respect to the standard inner-products, but they are orthogonal with respect to some new inner-products involving differential or difference operators. These classes of polynomials (called Sobolev orthogonal polynomials) have been the object of much attention in the last years (see [4], [11], [12], [14], [18], [25], [27], [28], [29], [30], [33], [37], [39], [41], [44], [45], [46], [47], [49], [52]).

2 The polynomials $\Psi_n(x)$

Definition 2 *A function $\mu(x):R\to R$ is called a distribution function if:*

1. *$\mu(x)$ is non-decreasing.*
2. *$\mu(x)$ is bounded.*
3. *$\mu(x)$ has finite moments, i.e.,* $\int\limits_{-\infty}^{\infty}x^n d\mu<\infty,\quad n\geq 0.$

The set

$$\mathfrak{S}\left(\mu\right)=\left\{x\in\mathbb{R}\mid\mu\left(x^{+}\right)-\mu\left(x^{-}\right)>0\right\}$$

is called the spectrum of $\mu(x)$.

We remind the reader of the following result, a proof of which can be found in [59].

Theorem 3 *A set of orthogonal polynomials* $\{p_n(x) \mid n \geq 0\}$ *satisfies a three term recurrence relation of the form*

$$p_{n+1} = (A_n x + B_n)\, p_n - C_n p_{n-1}, \quad n \geq 0 \tag{5}$$

with $p_{-1}(x) = 0$ *and* $p_0(x) = 1$.

The converse is known as Favard's Theorem [24].

Theorem 4 *Let* $\{p_n(x) \mid n \geq 0\}$ *be a sequence of polynomials, satisfying (5) with* $A_n \neq 0$ *for all* $n \geq 0$. *Then,*

(a) *There exists a function* $\mu(x)$ *of bounded variation on* $(-\infty, \infty)$ *such that*

$$\int_{-\infty}^{\infty} p_n(x) p_m(x) d\mu = M_n \delta_{n,m}, \quad n, m \geq 0$$

where

$$M_n = \mu_0 \frac{A_n}{A_0} \prod_{k=1}^{n} C_k$$

and $\mu_0 > 0$ *is a normalization constant.*

(b) $\mu(x)$ *can be chosen to be real valued if and only if* A_n, B_n *and* $C_n \in \mathbb{R}$ *for all* $n \geq 0$.

(c) $\mu(x)$ *is a distribution with an infinite spectrum if and only if* A_n, B_n *and* $C_n \in \mathbb{R}$ *for all* $n \geq 0$ *and* $M_n > 0$ *for all* $n \geq 1$.

Proof. See [17] for a proof and [1], [13], [22], [42], for some generalizations. ■

Therefore, if $\{\Psi_n(x) \mid n \geq 0\}$ is to be an orthogonal set, the polynomials $\Psi_n(x)$ must satisfy a recurrence relation of the form

$$\Psi_{n+1} = (A_n x + B_n)\, \Psi_n - C_n \Psi_{n-1} \tag{6}$$

with $\Psi_{-1}(x) = 0, \ \Psi_0(x) = 1$.

Let's define the function

$$G(x,z)=\exp\left[xf\left(z\right)\right]. \tag{7}$$

From (4) we see that in order to have $\Psi_0(x)=1$, we need $f(0)=0$. Also, from

$$\frac{\partial G}{\partial z}=xf'\left(z\right)G(x,z)=\sum_{n=0}^{\infty}\Psi_{n+1}(x)\frac{z^n}{n!}, \tag{8}$$

we have $f'\left(0\right)x=\Psi_1(x)$. If $\Psi_1(x)$ is to be a polynomial of degree 1, we need $f'\left(0\right)\neq 0$.

Taking the n^{th} derivative with respect to z in (8) and using Leibniz's rule, we get

$$\frac{\partial^{n+1}G}{\partial z^{n+1}}(x,z)=x\sum_{k=0}^{n}\binom{n}{k}f^{(n+1-k)}(z)\frac{\partial^k G}{\partial z^k}(x,z),\quad n\geq 0. \tag{9}$$

Setting $z=0$ in (9) and using (4) we obtain

$$\begin{aligned}\Psi_{n+1}(x)&=x\sum_{k=0}^{n}\binom{n}{k}f^{(n+1-k)}(0)\Psi_k(x)\\&=xf'\left(0\right)\Psi_n(x)+x\sum_{k=0}^{n-1}\binom{n}{k}f^{(n+1-k)}(0)\Psi_k(x).\end{aligned} \tag{10}$$

Comparing (10) with (6) we conclude that

$$A_n=f'\left(0\right)\equiv a \tag{11}$$

and

$$B_n\Psi_n-C_n\Psi_{n-1}=x\sum_{k=0}^{n-1}\binom{n}{k}f^{(n+1-k)}(0)\Psi_k(x). \tag{12}$$

Using (10) in (12) we have

$$\begin{aligned}&B_nx\sum_{k=0}^{n-1}\binom{n-1}{k}f^{(n-k)}(0)\Psi_k(x)-C_nx\sum_{k=0}^{n-2}\binom{n-2}{k}f^{(n-k-1)}(0)\Psi_k(x)\\&=x\sum_{k=0}^{n-1}\binom{n}{k}f^{(n+1-k)}(0)\Psi_k(x),\end{aligned} \tag{13}$$

which implies

$$B_n f'(0)\,\Psi_{n-1}(x) = n f''(0)\,\Psi_{n-1}(x)$$

or

$$B_n = n\frac{f''(0)}{f'(0)} \equiv nb. \tag{14}$$

Rearranging terms in (13) and using (14), we get

$$\sum_{k=0}^{n-2}\left[nb\binom{n-1}{k}f^{(n-k)}(0) - C_n\binom{n-2}{k}f^{(n-k-1)}(0) - \binom{n}{k}f^{(n+1-k)}(0)\right]\Psi_k(x) = 0$$

which gives

$$\frac{-C_n}{n(n-1)} = \frac{f^{(k+1)}(0) - kbf^{(k)}(0)}{k(k-1)\,f^{(k-1)}(0)}. \tag{15}$$

For equation (15) to be valid for $2 \le k \le n,\ n \ge 2$, it is necessary that both sides be equal to the same constant $-c$, with $c > 0$. Thus,

$$\frac{-C_n}{n(n-1)} = -c$$

or

$$C_n = cn(n-1)\,, \quad c > 0 \tag{16}$$

and

$$f^{(k+1)}(0) - kbf^{(k)}(0) + ck(k-1)\,f^{(k-1)}(0) = 0, \quad k \ge 1. \tag{17}$$

The solution of the recurrence (17), with $f^{(1)}(0) = a$, is given by

$$f^{(k)}(0) = \frac{ac}{R^-\left(bR^+ - 2c\right)}\Gamma(k)\left[\left(R^+\right)^k - \left(R^-\right)^k\right], \quad k \ge 1 \tag{18}$$

where $\Gamma(\cdot)$ is the Gamma function and

$$R^\pm = \frac{1}{2}\left[b \pm \sqrt{b^2 - 4c}\right]. \tag{19}$$

From (18) and $f(0) = 0$, we get

$$f(z) = \frac{a}{\sqrt{b^2 - 4c}}\ln\left[\frac{1 - zR^-}{1 - zR^+}\right] \tag{20}$$

or

$$f(z) = \frac{2a}{\sqrt{b^2 - 4c}}\left[\operatorname{arctanh}\left(\frac{b - 2cz}{\sqrt{b^2 - 4c}}\right) - \operatorname{arctanh}\left(\frac{b}{\sqrt{b^2 - 4c}}\right)\right].$$

We summarize the results of this section in the following theorem.

Theorem 5 *If the family of polynomials* $\{\Psi_n(x) \mid n \geq 0\}$ *defined by (4) satisfies (6), then*

$$f(z) = \frac{2a}{\sqrt{b^2 - 4c}} \left[\operatorname{arctanh}\left(\frac{b - 2cz}{\sqrt{b^2 - 4c}} \right) - \operatorname{arctanh}\left(\frac{b}{\sqrt{b^2 - 4c}} \right) \right], \tag{21}$$

and

$$A_n = a, \quad B_n = nb, \quad C_n = cn\,(n-1)$$

with $a \neq 0$ *and* $c > 0$. *The recurrence relation (6) takes the form*

$$\Psi_{n+1} = (ax + nb)\,\Psi_n - cn\,(n-1)\,\Psi_{n-1}, \quad n \geq 0. \tag{22}$$

Remark 6 *In the remainder of the paper we shall stress the dependence of* $\Psi_n(x)$ *on* a, b *and* c *by writing* $\Psi_n(x) = \Psi_n(x; a, b, c)$. *Note that from (21) it follows that*

$$T_n(x) = \Psi_n(x; 1, 0, 1). \tag{23}$$

Remark 7 *Since* $C_1 = 0$, *it follows from Favard's theorem that the full set* $\{\Psi_n(x) \mid n \geq 0\}$ *is not orthogonal with respect to an inner-product generated by a distribution function. However, we shall see in the next section that* $\{\Psi_n(x) \mid n \geq 1\}$ *is.*

3 Representation of Ψ_n

In order to find a representation of $\Psi_n(x; a, b, c)$, in terms of the classic hypergeometric polynomials, we shall consider three different cases, depending on the sign of the discriminant in (19).

3.1 Case 1: $b^2 - 4c = 0,\ b \neq 0$

From (19), we have $R^+ = R^- = \frac{1}{2}b$. Taking the limit as $c \to \frac{b^4}{4}$ in (20) gives

$$f(z) = -\frac{2a}{b} \frac{\frac{b}{2}z}{\frac{b}{2}z - 1}. \tag{24}$$

The generating function for the Laguerre polynomials is given by [9]

$$(1 - z)^{-a-1} \exp\left[x \frac{z}{z - 1} \right] = \sum_{n=0}^{\infty} L_n^{(\alpha)}(x) z^n, \quad \alpha > -1. \tag{25}$$

Hence, from (24) and (25), we obtain

$$\Psi_n\left(x; a, b, \frac{b^4}{4}\right) = n!\left(\frac{b}{2}\right)^n \lim_{\alpha\to-1} L_n^{(\alpha)}\left(-\frac{2a}{b}x\right). \tag{26}$$

To find the limit in (26) we prove the following lemma.

Lemma 8 *Let ${}_rF_s$ be the hypergeometric function [40]. Then,*

$$\lim_{\omega\to 0}(\omega)_n \ {}_rF_s\left(\begin{matrix} a_1(\omega),\ldots,a_r(\omega) \\ \omega, b_2(\omega),\ldots,b_s(\omega)\end{matrix}\,\middle|\, x\right) \tag{27}$$

$$= x\Gamma(n)\frac{\displaystyle\prod_{j=1}^{r} a_j(0)}{\displaystyle\prod_{j=2}^{s} b_j(0)}\ {}_rF_s\left(\begin{matrix} a_1(0)+1,\ldots,a_r(0)+1 \\ 2, b_2(0)+1,\ldots,b_s(0)+1\end{matrix}\,\middle|\, x\right)$$

Proof.

$$\lim_{\omega\to 0}(\omega)_n \ {}_rF_s\left(\begin{matrix} a_1(\omega),\ldots,a_r(\omega) \\ \omega, b_2(\omega),\ldots,b_s(\omega)\end{matrix}\,\middle|\, x\right)$$

$$= \lim_{\omega\to 0}(\omega)_n \sum_{k=0}^{\infty}\frac{(a_1(\omega))_k\times\cdots\times(a_r(\omega))_k}{(\omega)_k(b_2(\omega))_k\times\cdots\times(b_s(\omega))_k(1)_k}x^k$$

$$= \lim_{\omega\to 0}\Gamma(\omega+n)\sum_{k=0}^{\infty}\frac{(a_1(\omega))_k\times\cdots\times(a_r(\omega))_k}{\Gamma(\omega+k)(b_2(\omega))_k\times\cdots\times(b_s(\omega))_k(1)_k}x^k$$

$$= \Gamma(n)\sum_{k=1}^{\infty}\frac{(a_1(0))_k\times\cdots\times(a_r(0))_k}{\Gamma(k)(b_2(0))_k\times\cdots\times(b_s(0))_k(1)_k}x^k$$

$$= \Gamma(n)\sum_{k=0}^{\infty}\frac{(a_1(0))_{k+1}\times\cdots\times(a_r(0))_{k+1}}{\Gamma(k+1)(b_2(0))_{k+1}\times\cdots\times(b_s(0))_{k+1}(1)_{k+1}}x^{k+1}$$

$$= x\Gamma(n)\frac{\displaystyle\prod_{j=1}^{r} a_j(0)}{\displaystyle\prod_{j=2}^{s} b_j(0)}\sum_{k=0}^{\infty}\frac{(a_1(0)+1)_k\times\cdots\times(a_r(0)+1)_k}{(1)_k(b_2(0)+1)_k\times\cdots\times(b_s(0)+1)_k(2)_k}x^k$$

$$= x\Gamma(n)\frac{\prod_{j=1}^{r} a_j(0)}{\prod_{j=2}^{s} b_j(0)}\ {}_rF_s\left(\begin{array}{c} a_1(0)+1,\ldots,a_r(0)+1 \\ 2, b_2(0)+1,\ldots,b_s(0)+1 \end{array}\middle|\, x\right)$$

■

Corollary 9 *We can extend the class of Laguerre polynomials by defining*

$$L_0^{(-1)} = 1, \quad L_n^{(-1)}(x) = -\frac{x}{n} L_{n-1}^{(1)}(x), \quad n \geq 1. \tag{28}$$

Proof. Using the definition [36]

$$L_n^{(\alpha)}(x) = \frac{(\alpha+1)_n}{n!}\ {}_1F_1\left(\begin{array}{c} -n \\ \alpha+1 \end{array}\middle|\, x\right), \tag{29}$$

and (27), we get

$$\lim_{\alpha\to-1} L_n^{(\alpha)}(x) = \frac{1}{n!} x\Gamma(n)(-n)\ {}_1F_1\left(\begin{array}{c} -n+1 \\ 2 \end{array}\middle|\, x\right).$$

From (29) we have

$${}_1F_1\left(\begin{array}{c} -n+1 \\ 2 \end{array}\middle|\, x\right) = \frac{(n-1)!}{(2)_{n-1}} L_{n-1}^{(1)}(x) = \frac{1}{n} L_{n-1}^{(1)}(x), \quad n \geq 1.$$

Therefore,

$$\lim_{\alpha\to-1} L_n^{(\alpha)}(x) = -\frac{x}{n} L_{n-1}^{(1)}(x), \quad n \geq 1.$$

■

Corollary 10 *We have the representation*

$$\Psi_n\left(x; a, b, \frac{b^4}{4}\right) = a\left(\frac{b}{2}\right)^{n-1} (n-1)! L_{n-1}^{(1)}\left(-\frac{2a}{b}x\right), \quad n \geq 1.$$

Remark 11 *If we define the inner product*

$$\langle f, g\rangle_\alpha = \int_{-\infty}^{\infty} f(x) g(x) e^{-x} x^\alpha \chi_{[0,\infty)}(x) dx, \tag{30}$$

we have [36]

$$\left\langle L_n^{(-1)}, L_m^{(-1)} \right\rangle_{-1} = \left\langle -\frac{x}{n} L_{n-1}^{(1)}, -\frac{x}{m} L_{m-1}^{(1)} \right\rangle_{-1}$$

$$= \frac{1}{nm} \int_0^\infty L_{n-1}^{(1)}(x) L_{m-1}^{(1)}(x) x^2 e^{-x} x^{-1} dx = \frac{1}{nm} \left\langle L_{n-1}^{(1)}, L_{m-1}^{(1)} \right\rangle_1$$

$$= \frac{1}{n^2} \frac{\Gamma(n+1)}{(n-1)!} \delta_{n-1,m-1} = \frac{1}{n} \delta_{n,m}, \quad n, m \geq 1.$$

Therefore, $\left\{ L_n^{(-1)} \mid n \geq 1 \right\}$ *is an orthogonal set with respect to the inner product* $\langle \cdot, \cdot \rangle_{-1}$ *defined by (30). The general case was studied in [23] and [34], where it was shown that* $\left\{ L_n^{(-k)} \mid n \geq k \right\}$ *is an orthogonal set with respect to the inner product* $\langle \cdot, \cdot \rangle_{-k}$ *for all* $k \geq 1$.

Remark 12 *We will now show that* $\left\{ L_n^{(-1)} \mid n \geq 0 \right\}$ *is an orthonormal set with respect to the inner product*

$$\langle f, g \rangle = f(0)g(0) + \int_0^\infty f'(x) g'(x) e^{-x} dx.$$

From (28) we have

$$L_n^{(-1)}(0) = \delta_{n,0}, \quad n \geq 0, \tag{31}$$

which implies

$$\left\langle L_0^{(-1)}, L_n^{(-1)} \right\rangle = L_n^{(-1)}(0) + \int_0^\infty 0 e^{-x} dx = \delta_{n,0}, \quad n \geq 0.$$

Using the formula [40]

$$\frac{d}{dx} L_n^{(\alpha)} = -L_{n-1}^{(\alpha+1)}, \quad n \geq 0$$

and (31), we get

$$\left\langle L_n^{(-1)}, L_m^{(-1)} \right\rangle = L_n^{(-1)}(0) L_m^{(-1)}(0) + \int_0^\infty L_{n-1}^{(0)}(x) L_{m-1}^{(0)}(x) e^{-x} dx \tag{32}$$

$$= \delta_{n-1,m-1} = \delta_{n,m}, \quad n, m \geq 1.$$

Thus, (31) and (32) give

$$\left\langle L_n^{(-1)}, L_m^{(-1)} \right\rangle = \delta_{n,m}, \quad n, m \geq 0.$$

The general case for $L_n^{(-k)}$ was first considered in [38] and subsequently in [35], [43], [48], [50], [51], [53] and [60].

3.2 Case 2: $b^2 - 4c < 0$

From (19) we see that in this case $R^{\pm}$ are complex conjugates,

$$R^{\pm} = \frac{1}{2}\left[b \pm \mathrm{i}\sqrt{4c - b^2}\right] \tag{33}$$

with absolute value $|R^{\pm}| = \sqrt{c}$. Hence, we have

$$\frac{R^{\pm}}{\sqrt{c}} = e^{\mathrm{i}\phi}, \quad 0 < \phi < \pi. \tag{34}$$

Using (34) in (20) we can write $f(z)$ as

$$f(z) = \frac{a\mathrm{i}}{\sqrt{4c - b^2}} \ln\left[\frac{1 - z\sqrt{c}e^{\mathrm{i}\phi}}{1 - z\sqrt{c}e^{-\mathrm{i}\phi}}\right]. \tag{35}$$

Comparing (35) and the generating function for the Meixner-Pollaczek polynomials, given by

$$\left(1 - ze^{\mathrm{i}\phi}\right)^{-\lambda + \mathrm{i}x} \left(1 - ze^{-\mathrm{i}\phi}\right)^{-\lambda - \mathrm{i}x} = \sum_{n=0}^{\infty} P_n^{(\lambda)}(x; \phi) z^n, \quad \lambda > 0, \quad 0 < \phi < \pi, \tag{36}$$

we conclude that

$$\Psi_n(x; a, b, c) = n! c^{\frac{n}{2}} \lim_{\lambda \to 0} P_n^{(\lambda)}\left(\frac{ax}{\sqrt{4c - b^2}}; \phi\right), \quad b^2 - 4c < 0 \tag{37}$$

with ϕ defined by (34). We will then find the limit in (37) using Lemma 8.

Proposition 13 *The family of Meixner-Pollaczek polynomials can be extended if we define*

$$P_0^{(0)}(x; \phi) = 1, \quad P_n^{(0)}(x; \phi) = \frac{2x}{n} \sin(\phi) P_{n-1}^{(1)}(x; \phi), \quad n \geq 1. \tag{38}$$

Proof. Using the definition [36]

$$P_n^{(\lambda)}(x;\phi) = \frac{(2\lambda)_n}{n!}\ e^{\mathrm{i}n\phi}\ {}_2F_1\left(\begin{matrix} -n,\ \lambda+\mathrm{i}x \\ 2\lambda \end{matrix} \middle| 1-e^{-2\mathrm{i}\phi} \right) \tag{39}$$

and (27), we get

$$\begin{aligned} \lim_{\lambda\to 0} P_n^{(\lambda)}(x;\phi) &= \frac{1}{n!}\ e^{\mathrm{i}n\phi}\left(1-e^{-2\mathrm{i}\phi}\right)\Gamma(n)(-n)(\mathrm{i}x) \\ &\quad \times\ {}_2F_1\left(\begin{matrix} -n+1,\ \mathrm{i}x+1 \\ 2 \end{matrix} \middle| 1-e^{-2\mathrm{i}\phi} \right) \\ &= \frac{1}{n!}\ e^{\mathrm{i}n\phi}\left(1-e^{-2\mathrm{i}\phi}\right)\Gamma(n)(-n)(\mathrm{i}x)\,P_{n-1}^{(1)}(x;\phi)\frac{(n-1)!}{(2)_{n-1}}e^{-\mathrm{i}(n-1)\phi} \\ &= \frac{1}{n}\left(\frac{e^{\mathrm{i}\phi}-e^{-\mathrm{i}\phi}}{\mathrm{i}}\right) x P_{n-1}^{(1)}(x;\phi), \quad n\geq 1 \end{aligned}$$

and the result follows. ■

Corollary 14 *If* $b^2-4c<0$, *we have the representation*

$$\Psi_n(x;a,b,c) = (n-1)!c^{\frac{n-1}{2}}axP_{n-1}^{(1)}\left(\frac{a}{\sqrt{4c-b^2}}x;\phi\right), \quad n\geq 1 \tag{40}$$

with ϕ *defined by (34).*

Proof. Using (38) in (37) we have

$$\Psi_n(x;a,b,c) = n!c^{\frac{n}{2}}\frac{2x}{n}\frac{a}{\sqrt{4c-b^2}}\sin(\phi)\,P_{n-1}^{(1)}\left(\frac{a}{\sqrt{4c-b^2}}x;\phi\right), \quad n\geq 1.$$

From (33) and (34) we have

$$\frac{1}{2\sqrt{c}}\left[b\pm \mathrm{i}\sqrt{4c-b^2}\right] = \cos(\phi)\pm \mathrm{i}\sin(\phi),$$

which gives

$$\sin(\phi) = \sqrt{1-\frac{b^2}{4c}}.$$

■

Remark 15 *Replacing* $a = 1, b = 0$ *and* $c = 1$ *in (40) and using (23), we obtain*

$$T_n(x) = \Psi_n(x; 1, 0, 1) = (n-1)!xP_{n-1}^{(1)}\left(\frac{x}{2}; \frac{\pi}{2}\right), \quad n \geq 1$$

which agrees with (3).

Remark 16 *If we define the inner product*

$$\langle f, g \rangle_\lambda = \frac{1}{2\pi} \int\limits_{-\infty}^{\infty} f(x)g(x)e^{(2\phi-\pi)x} \left|\Gamma\left(\lambda + \mathrm{i}x\right)\right|^2 dx, \tag{41}$$

we have [36]

$$\begin{gathered}
\left\langle P_n^{(0)}, P_m^{(0)} \right\rangle_0 = \left\langle \frac{2x}{n} \sin\left(\phi\right) P_{n-1}^{(1)}\left(x; \phi\right), \frac{2x}{m} \sin\left(\phi\right) P_{m-1}^{(1)}\left(x; \phi\right) \right\rangle_0 \\
= \frac{4\sin^2\left(\phi\right)}{nm} \frac{1}{2\pi} \int\limits_{-\infty}^{\infty} P_{n-1}^{(1)}\left(x; \phi\right) P_{m-1}^{(1)}\left(x; \phi\right) e^{(2\phi-\pi)x} \left|\Gamma\left(\mathrm{i}x\right)\right|^2 x^2 dx \\
= \frac{4\sin^2\left(\phi\right)}{nm} \frac{1}{2\pi} \int\limits_{-\infty}^{\infty} P_{n-1}^{(1)}\left(x; \phi\right) P_{m-1}^{(1)}\left(x; \phi\right) e^{(2\phi-\pi)x} \left|\Gamma\left(1 + \mathrm{i}x\right)\right|^2 dx \\
= \frac{4\sin^2\left(\phi\right)}{nm} \left\langle P_{n-1}^{(1)}, P_{m-1}^{(1)} \right\rangle_1 \\
= \frac{4\sin^2\left(\phi\right)}{n^2} \frac{\Gamma\left(n+1\right)}{4\sin^2\left(\phi\right)\left(n-1\right)!} \delta_{n-1,m-1} = \frac{1}{n}\delta_{n,m}, \quad n, m \geq 1.
\end{gathered}$$

Therefore, $\left\{P_n^{(0)} \mid n \geq 1\right\}$ *is an orthogonal set with respect to the inner product* $\langle \cdot, \cdot \rangle_0$ *defined by (41).*

Remark 17 *We will now show that* $\left\{P_n^{(0)} \mid n \geq 0\right\}$ *is an orthonormal set with respect to the inner product*

$$\langle f, g \rangle = f(0)g(0) + \frac{1}{4\pi\sin\left(\phi\right)} \int\limits_{-\infty}^{\infty} \delta f(x)\delta g(x)e^{(2\phi-\pi)x} \left|\Gamma\left(\frac{1}{2} + \mathrm{i}x\right)\right|^2 dx,$$

where

$$\delta f(x) = f\left(x+\frac{1}{2}\right) - f\left(x-\frac{1}{2}\right).$$

From (38) we have

$$P_n^{(0)}(0;\phi) = \delta_{n,0}, \quad n \geq 0, \tag{42}$$

which implies

$$\left\langle P_0^{(0)}, P_n^{(0)} \right\rangle = P_n^{(0)}(0;\phi) + \int_{-\infty}^{\infty} 0 e^{(2\phi-\pi)x} \left|\Gamma\left(\frac{1}{2}+\mathrm{i}x\right)\right|^2 dx = \delta_{n,0}, \quad n \geq 0. \tag{43}$$

Using the formula [36]

$$\delta P_n^{(\lambda)}(x;\phi) = 2\sin(\phi)\, P_{n-1}^{\left(\lambda+\frac{1}{2}\right)}(x;\phi), \quad n \geq 0$$

and (42), we get

$$\begin{aligned} \left\langle P_n^{(0)}, P_m^{(0)} \right\rangle &= P_n^{(0)}(0;\phi) P_m^{(0)}(0;\phi) \\ &+ 2\sin(\phi)\,\frac{1}{2\pi}\int_{-\infty}^{\infty} P_{n-1}^{\left(\frac{1}{2}\right)}(x;\phi) P_{m-1}^{\left(\frac{1}{2}\right)}(x;\phi) e^{(2\phi-\pi)x} \left|\Gamma\left(\frac{1}{2}+\mathrm{i}x\right)\right|^2 dx \\ &= 2\sin(\phi)\,\frac{\Gamma(n)}{2\sin(\phi)\,(n-1)!}\delta_{n-1,m-1} = \delta_{n,m}, \quad n,m \geq 1. \end{aligned} \tag{44}$$

Thus, (43) and (44) give

$$\left\langle P_n^{(0)}, P_m^{(0)} \right\rangle = \delta_{n,m}, \quad n,m \geq 0.$$

3.3 Case 3: $b^2 - 4c > 0$

From (19) we have $R^- < R^+$, if $b^2 - 4c > 0$. Since $\sqrt{b^2-4c} < |b|$,

$$0 < R^- < R^+ \text{ for } b > 0 \text{ and } R^- < R^+ < 0 \text{ for } b < 0.$$

Thus,

$$0 < \frac{R^-}{R^+} < 1 \text{ for } b > 0 \text{ and } 0 < \frac{R^+}{R^-} < 1 \text{ for } b < 0. \tag{45}$$

Comparing (20) and the generating function for the Meixner polynomials, given by [54]

$$\left(1-\frac{t}{\gamma}\right)^{x}(1-t)^{-\beta-x}=\sum_{n=0}^{\infty}M_{n}^{(\beta)}(x;\gamma)\frac{t^{n}}{n!},\quad \beta>0,\quad 0<\gamma<1,$$

we have

$$\Psi_{n}\left(x;a,b,c\right)=\begin{cases}\left(R^{-}\right)^{n}\lim\limits_{\beta\to0}M_{n}^{(\beta)}\left(-\frac{ax}{\sqrt{b^{2}-4c}};\frac{R^{-}}{R^{+}}\right), & b>0\\ \left(R^{+}\right)^{n}\lim\limits_{\beta\to0}M_{n}^{(\beta)}\left(\frac{ax}{\sqrt{b^{2}-4c}};\frac{R^{+}}{R^{-}}\right), & b<0\end{cases}\tag{46}$$

where we have used (45). To find the limit in (46) we use Lemma 1.

Proposition 18 *The family of Meixner polynomials can be extended if we define*

$$M_{0}^{(0)}(x;\gamma)=1,\quad M_{n}^{(0)}(x;\gamma)=\left(1-\frac{1}{\gamma}\right)xM_{n-1}^{(2)}(x-1;\gamma),\quad n\geq1.\tag{47}$$

Proof. Using the definition [54]

$$M_{n}^{(\beta)}(x;\gamma)=(\beta)_{n}\ {}_{2}F_{1}\left(\begin{matrix}-n,\ -x\\ \beta\end{matrix}\ \middle|\ 1-\frac{1}{\gamma}\right)$$

and (27), we get

$$\begin{aligned}\lim_{\beta\to0}M_{n}^{(\beta)}(x;\gamma)&=\left(1-\frac{1}{\gamma}\right)\Gamma\left(n\right)(-n)(-x)\ {}_{2}F_{1}\left(\begin{matrix}-n+1,\ -x+1\\ 2\end{matrix}\ \middle|\ 1-\frac{1}{\gamma}\right)\\ &=\left(1-\frac{1}{\gamma}\right)xM_{n-1}^{(2)}(x-1;\gamma),\quad n\geq1\end{aligned}$$

and the result follows. ■

Corollary 19 *If* $b^{2}-4c>0$, *we have the representation*

$$\Psi_{n}\left(x;a,b,c\right)=\begin{cases}ax\left(R^{-}\right)^{n-1}M_{n-1}^{(2)}\left(-\frac{ax}{\sqrt{b^{2}-4c}}-1;\frac{R^{-}}{R^{+}}\right), & b>0\\ ax\left(R^{+}\right)^{n-1}M_{n-1}^{(2)}\left(\frac{ax}{\sqrt{b^{2}-4c}}-1;\frac{R^{+}}{R^{-}}\right), & b<0\end{cases},\quad n\geq1.\tag{48}$$

Proof. Using (47) in (48), we have

$$\begin{aligned}\Psi_n(x;a,b,c) &= \left(R^-\right)^n \left(1-\frac{R^-}{R^+}\right)\left(-\frac{ax}{\sqrt{b^2-4c}}\right) M_{n-1}^{(2)}\left(-\frac{ax}{\sqrt{b^2-4c}}-1;\frac{R^-}{R^+}\right)\\ &= \left(R^-\right)^{n-1}\left(-\sqrt{b^2-4c}\right)\left(-\frac{ax}{\sqrt{b^2-4c}}\right) M_{n-1}^{(2)}\left(\frac{ax}{\sqrt{b^2-4c}}-1;\frac{R^+}{R^-}\right)\end{aligned}$$

for $b>0$ and

$$\begin{aligned}\Psi_n(x;a,b,c) &= \left(R^+\right)^n \left(1-\frac{R^-}{R^+}\right)\frac{ax}{\sqrt{b^2-4c}} M_{n-1}^{(2)}\left(\frac{ax}{\sqrt{b^2-4c}}-1;\frac{R^+}{R^-}\right)\\ &= \left(R^+\right)^{n-1}\sqrt{b^2-4c}\frac{ax}{\sqrt{b^2-4c}} M_{n-1}^{(2)}\left(\frac{ax}{\sqrt{b^2-4c}}-1;\frac{R^+}{R^-}\right)\end{aligned}$$

for $b<0$. ■

Remark 20 *If we define the inner product*

$$\langle f,g\rangle_\beta = \sum_{k=0}^{\infty} f(k)g(k)\gamma^k \frac{\Gamma(\beta+k)}{k!}, \tag{49}$$

we have [54]

$$\left\langle M_n^{(\beta)}, M_m^{(\beta)}\right\rangle_\beta = \frac{n!\Gamma(\beta+n)}{\gamma^n(1-\gamma)^\beta}\delta_{n,m}, \quad n,m\geq 0.$$

Therefore,

$$\begin{aligned}\left\langle M_n^{(0)}, M_m^{(0)}\right\rangle_0 &= \left\langle \left(1-\frac{1}{\gamma}\right) xM_{n-1}^{(2)}(x-1;\gamma), \left(1-\frac{1}{\gamma}\right) xM_{m-1}^{(2)}(x-1;\gamma)\right\rangle_0\\ &= \left(1-\frac{1}{\gamma}\right)^2 \sum_{k=1}^{\infty} k^2 M_{n-1}^{(2)}(k-1;\gamma)M_{m-1}^{(2)}(k-1;\gamma)\gamma^k\frac{\Gamma(k)}{k!}\\ &= \left(1-\frac{1}{\gamma}\right)^2 \sum_{k=0}^{\infty} M_{n-1}^{(2)}(k;\gamma)M_{m-1}^{(2)}(k;\gamma)\gamma^k(k+1)\\ &= \left(1-\frac{1}{\gamma}\right)^2 \left\langle M_{n-1}^{(2)}, M_{m-1}^{(2)}\right\rangle_2\\ &= \left(1-\frac{1}{\gamma}\right)^2 \frac{(n-1)!\Gamma(n+1)}{\gamma^{n-1}(1-\gamma)^2}\delta_{n-1,m-1} = \frac{(n-1)!n!}{\gamma^{n+1}}\delta_{n,m}, \quad n,m\geq 1.\end{aligned}$$

Hence, $\left\{M_n^{(0)} \mid n \geq 1\right\}$ *is an orthogonal set with respect to the inner product* $\langle \cdot, \cdot \rangle_0$ *defined by (49).*

Remark 21 *We will now show that* $\left\{M_n^{(0)} \mid n \geq 0\right\}$ *is an orthogonal set with respect to the inner product*

$$\langle f, g \rangle = f(0)g(0) + \sum_{k=0}^{\infty} \Delta f(k) \Delta g(k) \frac{\gamma^k}{1-\gamma},$$

where

$$\Delta f(x) = f(x+1) - f(x).$$

From (47) we have

$$M_n^{(0)}(0;\gamma) = \delta_{n,0}, \quad n \geq 0, \tag{50}$$

which implies

$$\left\langle M_0^{(0)}, M_n^{(0)} \right\rangle = M_n^{(0)}(0;\gamma) + \sum_{k=0}^{\infty} 0 \frac{\gamma^k}{1-\gamma} = \delta_{n,0}, \quad n \geq 0. \tag{51}$$

Using the formula [54]

$$\Delta M_n^{(\beta)}(x;\gamma) = n \left(1 - \frac{1}{\gamma}\right) M_{n-1}^{(\beta+1)}(x;\gamma), \quad n \geq 0$$

and (50), we get

$$\begin{aligned} \left\langle M_n^{(0)}, M_m^{(0)} \right\rangle &= M_n^{(0)}(0;\gamma) M_m^{(0)}(0;\gamma) \\ &+ \sum_{k=0}^{\infty} n \left(1 - \frac{1}{\gamma}\right) M_{n-1}^{(1)}(k;\gamma) m \left(1 - \frac{1}{\gamma}\right) M_{m-1}^{(1)}(k;\gamma) \frac{\gamma^k}{1-\gamma} \\ &= n^2 \left(1 - \frac{1}{\gamma}\right)^2 \frac{(n-1)!\Gamma(n)}{\gamma^{n-1}(1-\gamma)^2} \delta_{n-1,m-1} \\ &= \frac{1}{\gamma^n} (n!)^2 \delta_{n,m}, \quad n, m \geq 1. \end{aligned} \tag{52}$$

Thus, (51) and (52) give

$$\left\langle M_n^{(0)}, M_m^{(0)} \right\rangle = \frac{1}{\gamma^n} (n!)^2 \delta_{n,m}, \quad n, m \geq 0.$$

Some extensions using a similar inner-product were studied in [5], [7], [8] and [21].

References

[1] N. I. Akhiezer. *The classical moment problem and some related questions in analysis.* Translated by N. Kemmer. Hafner Publishing Co., New York, 1965.

[2] W. A. Al-Salam. Characterization theorems for orthogonal polynomials. In *Orthogonal polynomials (Columbus, OH, 1989)*, volume 294 of *NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci.*, pages 1–24. Kluwer Acad. Publ., Dordrecht, 1990.

[3] W. A. Al-Salam and A. Verma. Generalized Sheffer polynomials. *Duke Math. J.*, 37:361–365, 1970.

[4] M. Álvarez de Morales, T. E. Pérez, and M. A. Piñar. Sobolev orthogonality for the Gegenbauer polynomials $\{C_n^{(-N+1/2)}\}_{n\geq 0}$. *J. Comput. Appl. Math.*, 100(1):111–120, 1998.

[5] M. Álvarez de Morales, T. E. Pérez, M. A. Piñar, and A. Ronveaux. Non-standard orthogonality for Meixner polynomials. *Electron. Trans. Numer. Anal.*, 9:1–25 (electronic), 1999. Orthogonal polynomials: numerical and symbolic algorithms (Leganés, 1998).

[6] T. K. Araaya. The Meixner-Pollaczek polynomials and a system of orthogonal polynomials in a strip. *J. Comput. Appl. Math.*, 170(2):241–254, 2004.

[7] I. Area, E. Godoy, and F. Marcellán. Inner products involving differences: the Meixner-Sobolev polynomials. *J. Differ. Equations Appl.*, 6(1):1–31, 2000.

[8] I. Area, E. Godoy, F. Marcellán, and J. J. Moreno-Balcázar. Ratio and Plancherel-Rotach asymptotics for Meixner-Sobolev orthogonal polynomials. *J. Comput. Appl. Math.*, 116(1):63–75, 2000.

[9] R. Askey. *Orthogonal polynomials and special functions.* Society for Industrial and Applied Mathematics, Philadelphia, Pa., 1975.

[10] P. Barrucand. Sur les polynômes d'Appell généralisés. *C. R. Acad. Sci. Paris Sér. A-B*, 274:A1109–A1111, 1972.

[11] H. Bavinck and H. G. Meijer. Orthogonal polynomials with respect to a symmetric inner product involving derivatives. *Appl. Anal.*, 33(1-2):103–117, 1989.

[12] H. Bavinck and H. G. Meijer. On orthogonal polynomials with respect to an inner product involving derivatives: zeros and recurrence relations. *Indag. Math. (N.S.)*, 1(1):7–14, 1990.

[13] M. Bertola. Bilinear semiclassical moment functionals and their integral representation. *J. Approx. Theory*, 121(1):71–99, 2003.

[14] A. Cachafeiro, F. Marcellán, and J. J. Moreno-Balcázar. On asymptotic properties of Freud-Sobolev orthogonal polynomials. *J. Approx. Theory*, 125(1):26–41, 2003.

[15] L. Carlitz. Some arithmetic properties of a special sequence of polynomials. *Duke Math. J*, 26:583–590, 1959.

[16] L. Carlitz. Some arithmetic properties of a special sequence of polynomials in the Gaussian field. *Duke Math. J.*, 30:15–24, 1963.

[17] T. S. Chihara. *An introduction to orthogonal polynomials*. Gordon and Breach Science Publishers, New York, 1978. Mathematics and its Applications, Vol. 13.

[18] M. Á. De Morales, T. E. Pérez, and M. A. Piñar. Orthogonal polynomials associated with a Δ-Sobolev inner product. *J. Difference Equ. Appl.*, 8(2):125–151, 2002.

[19] A. Di Bucchianico. Representations of Sheffer polynomials. *Stud. Appl. Math.*, 93(1):1–14, 1994.

[20] A. Di Bucchianico and D. Loeb. A simpler characterization of Sheffer polynomials. *Stud. Appl. Math.*, 92(1):1–15, 1994.

[21] A. Draux and C. Elhami. On the positivity of some bilinear functionals for discrete Sobolev orthogonal polynomials. *Rend. Mat. Appl. (7)*, 21(1-4):39–55, 2001.

[22] A. J. Duran. A generalization of Favard's theorem for polynomials satisfying a recurrence relation. *J. Approx. Theory*, 74(1):83–109, 1993.

[23] W. N. Everitt, L. L. Littlejohn, and R. Wellman. The Sobolev orthogonality and spectral analysis of the Laguerre polynomials $\{L_n^{-k}\}$ for positive integers k. *J. Comput. Appl. Math.*, 171(1-2):199–234, 2004.

[24] J. Favard. Sur les polynomes de Tchebycheff. *C. R. Acad. Sci. Paris*, 200:2052–2053, 1935.

[25] S. S. Han, I. H. Jung, K. H. Kwon, and J. K. Lee. A non-standard class of Sobolev orthogonal polynomials. *Commun. Korean Math. Soc.*, 12(4):935–950, 1997.

[26] J. Hofbauer. A representation of Sheffer polynomials in terms of a differential equation for their generating functions. *Aequationes Math.*, 23(2-3):156–168, 1981.

[27] A. Iserles, P. E. Koch, S. P. Nørsett, and J. M. Sanz-Serna. On polynomials orthogonal with respect to certain Sobolev inner products. *J. Approx. Theory*, 65(2):151–175, 1991.

[28] H. S. Jung, K. H. Kwon, and D. W. Lee. Discrete Sobolev orthogonal polynomials and second order difference equations. *J. Korean Math. Soc.*, 36(2):381–402, 1999.

[29] I. H. Jung, K. H. Kwon, D. W. Lee, and L. L. Littlejohn. Differential equations and Sobolev orthogonality. In *Proceedings of the International Conference on Orthogonality, Moment Problems and Continued Fractions (Delft, 1994)*, volume 65, pages 173–180, 1995.

[30] I. H. Jung, K. H. Kwon, and G. J. Yoon. Differential equations of infinite order for Sobolev-type orthogonal polynomials. *J. Comput. Appl. Math.*, 78(2):277–293, 1997.

[31] S. Kaijser. Some "new" orthogonal polynomials. *Normat*, 47(4):156–165, 188, 1999.

[32] R. Kelisky. The numbers generated by exp(arctan x). *Duke Math. J*, 26:569–581, 1959.

[33] D. H. Kim, K. H. Kwon, F. Marcellán, and S. B. Park. Sobolev-type orthogonal polynomials and their zeros. *Rend. Mat. Appl. (7)*, 17(3):423–444, 1997.

[34] E. Kochneff. Expansions in Laguerre polynomials of negative order. *J. Approx. Theory*, 81(3):332–346, 1995.

[35] J. Koekoek, R. Koekoek, and H. Bavinck. On differential equations for Sobolev-type Laguerre polynomials. *Trans. Amer. Math. Soc.*, 350(1):347–393, 1998.

[36] R. Koekoek and R. F. Swarttouw. The Askey-scheme of hypergeometric orthogonal polynomials and its q-analogue. Technical Report 98-17, Delft University of Technology, 1998. http://aw.twi.tudelft.nl∼koekoek/askey/.

[37] K. H. Kwon, D. W. Lee, and L. L. Littlejohn. Sobolev orthogonal polynomials and second order differential equation. II. *Bull. Korean Math. Soc.*, 33(1):135–170, 1996.

[38] K. H. Kwon and L. L. Littlejohn. The orthogonality of the Laguerre polynomials $\{L_n^{-k}(x)\}$ for positive integers k. *Ann. Numer. Math.*, 2(1-4):289–303, 1995. Special functions (Torino, 1993).

[39] K. H. Kwon and L. L. Littlejohn. Sobolev orthogonal polynomials and second-order differential equations. *Rocky Mountain J. Math.*, 28(2):547–594, 1998.

[40] N. N. Lebedev. *Special functions and their applications*. Dover Publications Inc., New York, 1972. Revised edition, translated from the Russian and edited by Richard A. Silverman, Unabridged and corrected republication.

[41] F. Marcellán, M. Alfaro, and M. L. Rezola. Orthogonal polynomials on Sobolev spaces: old and new directions. In *Proceedings of the Seventh Spanish Symposium on Orthogonal Polynomials and Applications (VII SPOA) (Granada, 1991)*, volume 48, pages 113–131, 1993.

[42] F. Marcellán and R. Álvarez-Nodarse. On the "Favard theorem" and its extensions. *J. Comput. Appl. Math.*, 127(1-2):231–254, 2001. Numerical analysis 2000, Vol. V, Quadrature and orthogonal polynomials.

[43] F. Marcellán, H. G. Meijer, T. E. Pérez, and M. A. Piñar. An asymptotic result for Laguerre-Sobolev orthogonal polynomials. *J. Comput. Appl. Math.*, 87(1):87–94, 1997.

[44] F. Marcellán and L. Moral. Sobolev-type orthogonal polynomials on the unit circle. *Appl. Math. Comput.*, 128(2-3):329–363, 2002. Orthogonal systems and applications.

[45] F. Marcellán, T. E. Pérez, and M. A. Piñar. Gegenbauer-Sobolev orthogonal polynomials. In *Nonlinear numerical methods and rational approximation, II (Wilrijk, 1993)*, volume 296 of *Math. Appl.*, pages 71–82. Kluwer Acad. Publ., Dordrecht, 1994.

[46] F. Marcellán, T. E. Pérez, and M. A. Piñar. Orthogonal polynomials on weighted Sobolev spaces: the semiclassical case. *Ann. Numer. Math.*, 2(1-4):93–122, 1995. Special functions (Torino, 1993).

[47] F. Marcellán, T. E. Pérez, and M. A. Piñar. Regular Sobolev type orthogonal polynomials: the Bessel case. *Rocky Mountain J. Math.*, 25(4):1431–1457, 1995.

[48] F. Marcellán, T. E. Pérez, and M. A. Piñar. Laguerre-Sobolev orthogonal polynomials. *J. Comput. Appl. Math.*, 71(2):245–265, 1996.

[49] F. Marcellán, T. E. Pérez, M. A. Piñar, and A. Ronveaux. General Sobolev orthogonal polynomials. *J. Math. Anal. Appl.*, 200(3):614–634, 1996.

[50] F. Marcellán and J. Sánchez-Ruiz. Connection coefficients for Laguerre-Sobolev orthogonal polynomials. *J. Math. Anal. Appl.*, 283(2):440–458, 2003.

[51] H. G. Meijer. Laguerre polynomials generalized to a certain discrete Sobolev inner product space. *J. Approx. Theory*, 73(1):1–16, 1993.

[52] H. G. Meijer. A short history of orthogonal polynomials in a Sobolev space. I. The non-discrete case. *Nieuw Arch. Wisk. (4)*, 14(1):93–112, 1996. 31st Dutch Mathematical Conference (Groningen, 1995).

[53] H. G. Meijer and M. A. Piñar. A generating function for Laguerre-Sobolev orthogonal polynomials. *J. Approx. Theory*, 120(1):111–123, 2003.

[54] A. F. Nikiforov, S. K. Suslov, and V. B. Uvarov. *Classical orthogonal polynomials of a discrete variable.* Springer Series in Computational Physics. Springer-Verlag, Berlin, 1991. Translated from the Russian.

[55] E. C. Popa. Note on Sheffer polynomials. *Octogon Math. Mag.*, 5(2):56–57, 1997.

[56] E. C. Popa. On the Sheffer polynomials. *Bul. Ştiinţ. Univ. Politeh. Timiş. Ser. Mat. Fiz.*, 43(57)(1):21–23, 1998.

[57] I. M. Sheffer. Some properties of polynomial sets of type zero. *Duke Math. J.*, 5:590–622, 1939.

[58] I. M. Sheffer. Some applications of certain polynomial classes. *Bull. Amer. Math. Soc.*, 47:885–898, 1941.

[59] N. M. Temme. *Special functions*. A Wiley-Interscience Publication. John Wiley & Sons Inc., New York, 1996. An introduction to the classical functions of mathematical physics.

[60] J. Wimp and H. Kiesel. Some properties of the Sobolev-Laguerre polynomials. In *Proceedings of the VIIIth Symposium on Orthogonal Polynomials and Their Applications (Seville, 1997)*, volume 99, pages 401–412, 1998.